

# Solution of the Master Equation for Quantum Brownian Motion given by the Schrödinger Equation

R. Sinuvasan<sup>1</sup>, A Paliathanasis<sup>\*2,3</sup>, R.M Morris<sup>4</sup> and P.G.L. Leach<sup>†4,5</sup>

<sup>1</sup>Department of Mathematics, Pondicherry University, Kalapet Puducherry 605 014, India

<sup>2</sup>Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Valdivia, Chile

<sup>3</sup>Institute of Systems Science, Durban University of Technology, PO Box 1334, Durban 4000,  
Republic of South Africa

<sup>4</sup>Department of Mathematics and Institute of Systems Science, Research and Postgraduate  
Support, Durban University of Technology, PO Box 1334, Durban 4000, Republic of South  
Africa

<sup>5</sup>School of Mathematics, Statistics and Computer Science, University of KwaZulu-Natal,  
Private Bag X54001, Durban 4000, Republic of South Africa

December 14, 2016

## Abstract

We consider the master equation of quantum Brownian motion and with the application of the group invariant transformation we show that there exists a surface on which the solution of the master equation is given by an autonomous one-dimensional Schrödinger Equation.

**Keywords:** Quantum Brownian motion, Master equation, Group invariant transformations

## 1 Introduction

With the method of path integrals, and specifically the Feynman-Vernon influence functional [1], Haake and Reibold [2] and few years latter Hu, Paz and Zhang derived an equation which inherits the properties of quantum Brownian motion for a harmonic oscillator interacting with a linear passive heat bath of oscillators [3, 4]. An alternate derivation of that master equation has been performed by Halliwell and Yu by tracing the evolution equation for the Wigner function of the system [5].

The master equation of quantum Brownian motion is an  $(1 + 2)$  linear nonautonomous evolution equation given by

$$Z_{,t} = -\frac{x}{m}Z_{,y} + m\Omega^2(t)yZ_{,x} + 2\Gamma(t)(xZ)_{,x} + \hbar m\Gamma(t)h(t)Z_{,xx} + \hbar\Gamma(t)f(t)Z_{,xy}, \quad (1)$$

where  $m$  is the mass of the Brownian particle,  $Z = Z(t, x, y)$  is the Wigner function of the density matrix ( $x$  denotes the momentum of the oscillator and  $y$  its position). Furthermore, the coefficients,  $\Omega^2(t)$ ,  $\Gamma(t)$ ,  $h(t)$

---

<sup>\*</sup>anpaliat@phys.uoa.gr

<sup>†</sup>leach@ucy.ac.cy

and  $f(t)$ , in general are time-dependent and related to the natural frequency of the Brownian motion and the terms interacting with the heat bath of oscillators. The derivation of the coefficients is given in [5, 6].

A general analytical solution of the master equation (1) with the use of the Langevin Equation has been derived in [6], whereas in [7] some solutions of the master equations for quantum Brownian motions are given. The analysis of open quantum system does not stop in equation (1). A relation of the exact master equation with the nonequilibrium Green functions for non-Markovian open quantum systems was derived in [8], while new phenomena concerning the thermal-state of a quantum system were predicted for a strongly non-Markovian environment in [9].

In this work we are interested in the existence of solutions for equation (1) which follow from the method of group invariant transformations, in particular we are interested in the one-parameter point transformations which were introduced by S. Lie [10], where the generator of the infinitesimal transformation is called a Lie (point) symmetry. The importance of Lie symmetries is that they provide a systematic method to facilitate the solution of differential equations because they provide first-order invariants which can be used to reduce the order of differential equations. Moreover Lie symmetries can be used for the classification of differential equations and important information for the differential equation can be extracted from the admitted group of invariant transformations. The method of group invariant transformations has been applied in various systems of quantum mechanics, for instance see [11, 12, 13, 14, 15, 16] and references therein.

By applying the Lie theory for differential equations we show that equation (1) is invariant under a group of one-parameter point transformations in which the generators form the  $\{A_1 \oplus_s W_5\} \oplus_s \infty A_1$ , Lie algebra, where  $W_5$  denotes the five-element Weyl-Heisenberg algebra and  $\infty A_1$  is the infinite-dimensional abelian algebra of the solutions of the linear  $(1+2)$  evolution equation and follows from the linearity of (1). Furthermore from the Lie symmetries we can define a surface in which equation (1) is independent of one of the independent variables and with the use of the zeroth-order invariants we can reduce equation (1) in a nonautonomous one-dimensional evolution equation. We study the Lie point symmetries of this equation and show that is maximally symmetric. Hence it is invariant under a group of transformation which form the<sup>1</sup>  $\{sl(2, R) \oplus_s W_3\} \oplus_s \infty A_1$  Lie algebra. From S. Lie's theorem this indicates that there exists a "coordinate" transformation in which the reduced equation is equivalent to the equation. Hence solutions of the Schrödinger equation are also solutions of the master equation (1). The plan of the paper is as follows.

In Section 2 we give the basic properties and definitions of Lie symmetries and we study the existence of Lie symmetries for the master equation (1). Furthermore we apply the zeroth-order invariants of the Lie symmetries and we reduce the original equation to a one-dimensional evolution equation. In Section 3 we study the relationship between the reduced equation and the Schrödinger equation. Finally we draw our conclusions and give an example in Section 4.

## 2 Lie Point symmetries of the Master Equation

For the convenience of the reader we present the basic properties and definitions of Lie symmetries of differential equations.

Consider a differential equation  $\Theta(x^k, u, u_{,i}, u_{,ij}) = 0$ , where  $x^k$  are the independent variables, and  $u = u(x^k)$  is the dependent variable. Then the differential operator,

$$X = \xi^i(x^k, u) \partial_i + \eta(x^k, u) \partial_u, \quad (2)$$

is called a Lie symmetry of  $\Theta$ , if there exists a function  $\lambda$ , such that  $\mathcal{L}_{X^{[2]}} \Theta = \lambda \Theta$ , where  $X^{[2]}$  is the second prolongation/extension of the vector field  $X$ , in the space  $\{x^k, u, u_{,i}, u_{,ij}\}$  [21, 22].

---

<sup>1</sup>Algebra  $sl(2, R)$  is the  $A_{3,8}$  and  $W_3$  is the  $A_{3,3}$  in the Mubarakzyanov Classification Scheme [17, 18, 19, 20]

Lie symmetries of differential equations can be used in order to determine invariant solutions or transform solutions to solutions [22]. From the Lie symmetry condition one defines the associated Lagrange's system

$$\frac{dx^i}{\xi^i} = \frac{du}{\eta} = \frac{du_i}{\eta_{[i]}} = \dots = \frac{du_{ij\dots i_n}}{\eta_{[ij\dots i_n]}} \quad (3)$$

the solution of which provides the characteristic functions

$$\Lambda^{[0]}(x^k, u), \Lambda^{[1]i}(x^k, u, u_i), \dots, \Lambda^{[n]}(x^k, u, u_i, \dots, u_{ij\dots i_n}). \quad (4)$$

The solution  $\Lambda^{[k]}$  is called the  $k$ th-order invariant of the Lie symmetry vector, (2). These invariants can be used in order to reduce the order or the number of the independent variables of the differential equations. Another important feature of Lie symmetries of differential equations is that they span the Lie algebra  $G_L$ . The application of a Lie symmetry to  $\Theta$  leads to a new differential equation  $\bar{\Theta}$  which is different from  $\Theta$  and possibly admits Lie symmetries which are not Lie symmetries of  $\Theta$ . This means that the reduced equation can have properties different from the original equation. However, the solutions of these equations are related through the point transformation which transformed  $\Theta$ , to  $\bar{\Theta}$ .

## 2.1 The Master Equation

In order to simplify the presentation of the calculations we rewrite equation (1) in the following form<sup>2</sup>

$$-\frac{x}{m}Z_{,y} + p(t)yZ_{,x} + q(t)(xZ)_{,x} + r(t)Z_{,xx} + s(t)Z_{,xy} - Z_{,t} = 0, \quad (5)$$

where  $p(t) = m\Omega^2(t)$ ,  $q(t) = 2\Gamma(t)$ ,  $r(t) = \hbar m\Gamma(t)h(t)$ ,  $s(t) = \hbar\Gamma(t)f(t)$ .

We assume the generator of the one-paramete infinitesimalr point transformation to be

$$X = \xi^t \partial_t + \xi^x \partial_x + \xi^y \partial_y + \eta \partial_Z, \quad (6)$$

in which  $\xi^t, \xi^x, \xi^y$  and  $\eta$  are functions of  $\{t, x, y, Z\}$ . Furthermore, because equation (5) is a linear equation, we have that  $\eta = G(t, x, y)Z + G_0Z + b(t, x, y)$ , where  $b(t, x, y)$  are solutions of equation (5) and form the infinite-dimensional Lie algebra  $\infty A_1$ , [23].

Hence from the Lie symmetry condition we have that<sup>3</sup>

$$\xi^t = a(t), \quad \xi^y = f_1(t), \quad (7)$$

$$\xi^x = m \frac{-f_1 p s + 2r f_1' + m(q s f_1' + f_1' s' - s f_1'')}{(2r + m(2q s + s'))} \quad (8)$$

and

$$\begin{aligned} G = & -(2r + m(2q s + s'))^{-2} \\ & + s') + p2(x + myq)r + m2xqs + 2myq^2s \\ & + 2mysq' + 2yr' + xs' + 3myqs' + mys'' \\ & + m2m^2yq^3s f_1' - 2myr f_1' q' - m^2y f_1' q' s' \\ & + m q^2 f_1' (2yr + 2xs + 3mys') - y(p f_1' 2r + m(2qs \\ & + s')) + 2xr f_1'' + 2m^2ysq' f_1'' + 2myr' f_1'' + m x s' f_1'' + m^2y f_1'' s'' \\ & - 2myr f_1''' - m^2ys' f_1''' + q2xr f_1' + m(f_1' (2yr' + xs' + mys'')) \\ & + 2(xs f_1'' + mys' f_1'' - mys f_1'''))), \end{aligned} \quad (9)$$

<sup>2</sup>It is possible to apply also a coordinate transformation,  $(x, y) \rightarrow (\bar{x}, \bar{y})$ , which “diagonalises” the second derivatives in equation (1). However, we prefer to work on the original physical system.

<sup>3</sup>In this work we used the symbolic package Sym for Mathematica [24].

where  $G_0$  is a constant and prime means differentiation with respect to time, “ $t$ ”, and functions  $a(t)$  and  $f_1(t)$  are related to  $p, q, r, s$  by a system of ordinary differential equations which we omit. We can see that  $f_1(t)$  satisfies a linear fourth-order differential equation which means that it provides us with four symmetries. Another symmetry vector arises from the unique solution of  $a(t)$ . Therefore from (7)-(9) it is easy to see that the Lie symmetries of the master equation form the  $\{A_1 \oplus_s W_5\} \oplus_s \propto A_1$  Lie algebra.

Indeed the form of the symmetry vector (6) it is not a closed form. The reason for this is that we have considered arbitrary functions,  $\Omega^2(t)$ ,  $\Gamma(t)$ ,  $h(t)$  and  $f(t)$ . In a case for specific functional forms of the coefficients one can calculate the symmetry vector in closed-form. For instance in the case for which the coefficients,  $p, q, r$  and  $s$ , are constants the Lie symmetries are

$$Y_1 = a_1 \partial_t, \quad Y_Z = Z \partial_Z, \quad Y_b = b \partial_z, \quad (10)$$

$$X_1 = e^{\frac{\lambda-q}{2}t} [m(\lambda-q) \partial_x + 2\partial_y], \quad X_2 = e^{-\frac{\lambda+q}{2}t} [m(\lambda+q) \partial_x - 2\partial_y], \quad (11)$$

$$X_3 = e^{-\frac{(\lambda-q)}{2}t} [2rm(q-\lambda) \partial_x + 4(r+sqm) \partial_y + (2mq(\lambda-q)x + m^2q(\lambda^2 - q^2)y) Z \partial_Z] \quad (12)$$

and

$$X_4 = e^{\frac{\lambda+q}{2}t} [2rm(\lambda+q) \partial_x + (r+sqm) \partial_y + (-2mq(\lambda+q)x + m^2q(\lambda^2 - q^2)y) Z \partial_Z], \quad (13)$$

where  $\lambda = \sqrt{4p - mq^2}$ .

Below we apply the zeroth-order invariants of the Lie symmetry which corresponds to the solution of the function  $f_1(t)$ .

## 2.2 Application of the Lie invariants

Consider now the Lie symmetry vector (6) for which  $G_0 = 0$  and  $a(t) = 0$ . The characteristic functions are

$$Z(t, x, y) = U(t, B(t)f_1(t)x - A(t)y) \exp[J(t, x)(B(t)f_1(t)x - A(t)y)], \quad (14)$$

where

$$J = \frac{x}{2A^2} \left[ \begin{aligned} &A^2(2G + xH) + \\ &+ B(-2(Bf_1x - A(t)y) + xBf_1)K \end{aligned} \right] \quad (15)$$

and the functions,  $A(t)$ ,  $B(t)$ ,  $K(t)$ ,  $G(t)$  and  $H(t)$ , are the coefficients of the symmetry vector (6).

Hence the application of (14) to (5) gives the reduced equation

$$S(t)U_{,ww} - wR(t)U_{,w} + q(t)U - U_{,t} = 0, \quad (16)$$

where  $w = B(t)f_1(t)x - A(t)y$ ,  $S(t) = Bf_1(Bf_1r - As)$  and  $R(t) = A^{-1}(Bf_1p - A')$ .

This means that the Lie symmetries provide us with a solution for the master equation (1), which is (14) and the function  $U(t, w)$  is given by (16). Below we study the Lie symmetries of (16) and we show that it is maximally symmetric. This means that it is equivalent with the elementary one-dimensional Schrödinger Equation.

## 3 Equivalence with the Schrödinger Equation

For simplicity in the following we consider a “time” rescaling,  $t \rightarrow T$ , such that  $S(T) = 1$ . Hence equation (16) becomes

$$U_{,ww} - wR(T)U_{,w} + q(T)U - U_{,T} = 0. \quad (17)$$

We apply the Lie symmetry condition to the this equation and we derive the following symmetry vector field

$$Y = \alpha(T)\partial_t + \left[ \frac{\dot{\alpha}(T)v}{2} + \beta(T) \right] \partial_v + (F(T, w)U + \bar{b}(t, w)) \partial_U, \quad (18)$$

where

$$F(T, v) = \phi(T) + \frac{1}{4} \left[ 2v\beta(T)R(T) + v^2R(T)\dot{\alpha} - 2v\dot{\beta} + v^2\alpha(T)\dot{R} - \frac{1}{2}v^2\ddot{\alpha} \right], \quad (19)$$

in which overdot denotes differentiation with respect to  $T$  and the functions  $\phi(T)$ ,  $\beta(T)$  and  $\alpha(T)$  are solutions of the equations

$$\phi = \phi_0 + \alpha \left( q + \frac{1}{2}R \right) - \frac{1}{4}\dot{\alpha}, \quad (20)$$

$$\ddot{\beta} = (\dot{R} + R^2)\beta \quad \text{and} \quad (21)$$

$$\ddot{\alpha} = 4\dot{\alpha}(\dot{R} + R^2) + 2\alpha \frac{d}{dT}(\dot{R} + R^2). \quad (22)$$

Furthermore  $\bar{b}(t, w)$  satisfies the original equation, (17).

Equation (21) is a maximally symmetric linear second-order differential equation. In this case, by application of the Riccati transformation  $R = \frac{\dot{\beta}}{\beta}$ , in (21) we find the solution,

$$\beta(T) = \beta_0 L(T) + \beta_1 L(T) \int L^{-2}(T) dT. \quad (23)$$

Equation (22) is a nonautonomous third-order differential equation. We multiply with  $\alpha(T)$  and integrate to obtain

$$\alpha(T)\ddot{\alpha} - \frac{1}{2}\dot{\alpha}^2 - 2\alpha^2(T)(\dot{R} + R^2(T)) = 2K,$$

where  $K$  is a constant. We substitute  $\alpha = \gamma^2$  into this equation and hence we find the well-known Ermakov-Pinney equation [25, 26]

$$\ddot{\rho} - \rho(T)(\dot{R} + R^2(T)) = \frac{K}{\rho^3(T)}. \quad (24)$$

The solution of (24) is given in [26] and it is related with the solution of the linear equation

$$\ddot{\sigma} - (\dot{R} + R^2(T))\sigma = 0. \quad (25)$$

Therefore we conclude that equation (17) admits as Lie symmetries the vector fields which form the  $\{sl(2, R) \oplus_s W_3\} \oplus_s \infty A_1$  Lie algebra. Hence from S. Lie's theorem [10] we have that there exists a transformation,  $(T, w, U) \rightarrow (\tau, \chi, \Psi)$ , in which (17) becomes

$$-\frac{\hbar}{2M} \frac{\partial^2 \Psi}{\partial \chi^2} = i\hbar^2 \frac{\partial \Psi}{\partial \tau} \quad (26)$$

which is the Schrödinger equation for a free particle. That is possible because equations (17) and (26) are both maximally symmetric.

## 4 Discussion

In this work with the application of the group invariant transformations we proved that there exists a surface in the space of the dependent and independent variables in which the master equation (1) can be seen as a one-dimensional equation. That means that solutions of the latter generate solutions for the master equation

given by the expression (14), that is, there is class of solutions which describe the two-different systems, but the solutions are given in different representations.

We remark that in our analysis we considered that the coefficients of the master equation are arbitrary functions of time, which means that the result holds when the coefficients are constants. For instance, consider the application of the Lie symmetry  $X_3$ , (11), in equation (5) for constant coefficients. Hence we have that  $Z = U(t, w)$ , where  $w = \frac{ym(L-q)-2x}{m(L-q)}$ , and  $U(t, w)$  satisfies the equation

$$\bar{s}U_{,ww} - \frac{(\lambda + q)}{2}wU_{,w} + 2qU - U_{,t} = 0 \quad (27)$$

and  $\bar{s} = 2\frac{-2r+sm(\lambda-q)}{m^2(\lambda-q)^2}$ . Therefore under the coordinate transformation,

$$U(t, w) = e^{2qt}\Psi(\tau, \chi) \ , \ w = \left(\frac{2Ms}{\hbar}\right)^{1/2} \chi e^{\left(\frac{\lambda+q}{2}t\right)} \ , \ d\tau = -i\hbar e^{-(\lambda+q)t} dt, \quad (28)$$

the latter equation takes the form of the one-dimensional Schrödinger equation. A similar result holds and for the remaining Lie symmetry vectors,  $X_2 - X_4$ , or any linear combination of them.

## Acknowledgements

The research of AP was supported by FONDECYT postdoctoral grant no. 3160121. AP thanks the Durban University of Technology for the hospitality provided while part of this work was performed. RMM thanks the National Research Foundation of the Republic of South Africa for the granting of a postdoctoral fellowship with grant number 93183 while this work was being undertaken. RS thanks the University Grants Commission of India for support.

## References

- [1] R.P. Feynman and F.L. Vernov, *Annals of Physics*, **24** 118 (1963)
- [2] F. Haake and R. Reibold, *Phys. Rev. A* **32**, 2462 (1985)
- [3] B.L. Hu, J.P. Paz and Y. Zhang, *Phys. Rev. D*, **45** 2843 (1992)
- [4] B.L. Hu, J.P. Paz and Y. Zhang, *Phys. Rev. D*, **47** 1576 (1993)
- [5] J.J. Halliwell and T. Yu, *Phys. Rev. D*, **53** 2012 (1996)
- [6] G.W. Ford, R.F. O'Connell, *Phys. Rev. D*, **64** 105020 (2001)
- [7] C.H. Fleming, A. Roura and B.L. Hu, *Annals of Physics*, **326**, 1207 (2011)
- [8] W.-M. Zhang, P.-Y. Lo, H.-N. Xiong, M. Wei-Yuan and F. Nori, *Phys. Rev. Lett.* **109**, 170402 (2012)
- [9] H.-N. Xiong, P.-Y. Lo, W.-M. Zhang, D. H. Feng and F. Nori, *Scientific Reports* **5**, 13353 (2015)
- [10] S. Lie, *Lectures on differential equations with known infinitesimal transformations*, (in German, written with the help of G Scheffers), Teubner BG, Leipzig, (1891)
- [11] J. Belmonte-Beitia, V.M. Perez-Garcia, V. Vekslerchik and P.J. Torres, *Phys. Rev. Lett.*, **98** 064102 (2007)
- [12] L. Gagnon and P. Winternitz, *J. Phys. A: Math. Gen.* **21** 1493 (1988)

- [13] L. Gagnon and P. Winternitz, J. Phys. A: Math. Gen. **22** 469 (1988)
- [14] R.O. Popovych, N.M. Ivanova and H. Eshraghi, J. Math. Phys. **45** 3049 (2014)
- [15] A. Paliathanasis and M. Tsamparlis, Int. J. Geom. Meth. Mod. Phys. **11** 1450037 (2014)
- [16] M.B. Sheftel, P. Tempesta and P. Winternitz, Physics of Atomic Nuclei, **65** 1144 (2002)
- [17] V.V. Morozov, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika*, **5** 161 (1958)
- [18] G.M. Mubarakzyanov, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika*, **32** 114 (1963)
- [19] G.M. Mubarakzyanov, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika*, **34** 99 (1963)
- [20] G.M. Mubarakzyanov, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika*, **35** 104 (1963)
- [21] P.J. Olver, Applications of Lie Groups to Differential Equations, Graduate Texts in Mathematics, Volume **107**, Springer-Verlag, New York, (1993)
- [22] G.W Bluman and S. Kumei, Symmetries of Differential Equations, Springer-Verlag, New York, (1989)
- [23] G.W. Bluman, J. Math. Anal. Appl. **145** 52 (1990)
- [24] S. Dimas and D. Tsoubelis, SYM: A new symmetry-finding package for Mathematica, Group Analysis of Differential Equations, Ed. by N.H. Ibragimov, C. Sophocleous and P.A. Damianou, University of Cyprus, Nicosia, 64 (2005)
- [25] V.P. Ermakov, Univ. Isz. Kiev Ser. III **9** p1, translated by A.O. Harin (1880)
- [26] E. Pinney, Proc. Am. Math. Soc. **1** 681 (1950)